

Transdermal mesotherapy treatments

MACDERMOL[®]



Technical data

	MACDERMOL [®] Bio Revitalisation	MACDERMOL [®] Bio Lift	MACDERMOL [®] Body Contour	MACDERMOL [®] Scalp Stimulation
Composition	Hyaluronic acid (2 mg /ml) + specific complex	Hyaluronic acid (2 mg /ml) + specific complex	Hyaluronic acid (2 mg /ml) + specific complex	Hyaluronic acid (2 mg /ml) + specific complex
Indication	For patients with dehydrated, tired and dull skin	For patients with relaxed and atonic skin	Treatment of gynoid lipodystrophy (cellulite) or localized fat deposits	Treatment of hair loss
Effects	Revitalizes and provides a long-lasting hydration to prevent and combat skin ageing.	Provides an immediate lifting effect. Tightens and tones slack skin	Reduces the appearance of cellulite. Lipolysis	Slows down hair loss and stimulates hair growth
Recommended zones	Face, neck, décolleté, back of hands, of arms and thighs.	Face, neck, décolleté, abdomen, knee, internal face	Upper thighs, hips, buttocks, abdomen, back of arms.	Scalp
Administration	Transdermal delivery (electrophoresis, iontophoresis, ultrasonophoresis, electroporation, etc.).			
Packaging	10 x 10 ml multidose glass vials containing each 5 ml of solution			
Sterilization	Microfiltration			
Storage	Between +2°C and 25°C			
Shelf life	3 years			

Presentation

Novatex has developed four transdermal mesotherapy treatments presented as a box of 10x10 ml multidose glass vials containing each 5 ml of non-animal hyaluronic acid (2 mg/ml) and specific complex.

How to use

Injection by transdermal delivery (electrophoresis, iontophoresis, ultrasonophoresis, electroporation, etc.).

Envisage 2 to 3 sessions at intervals of 3 to 4 weeks according to the result, then a session approximately every 3 to 4 months for maintenance.



Manufacturer :

NOVATEX BIOENGINEERING SA
Schwarzwaldallee 215
CH-4002 Basel
Switzerland

www.novatex-bioeng.com